

## A quick note on orthogonality relation (supplement of Slides #6)

In Eq. (14) in Slides #6, we have

$$\sum_{n=1}^{\infty} a_n \sin(n\pi x) = F(x) \quad , \quad (14) \text{ of Slides \#6}$$

where  $F(x) = 4\sin(3\pi x) + 7\sin(8\pi x)$ . How do we determine the coefficients,  $a_n$ , using the given information? In this simple example, a quick observation suffices for us to conclude that

$$a_3 = 4, \quad a_8 = 7, \quad \text{and } a_n = 0 \text{ for all other } n .$$

However, if the  $F(x)$  has a more complicated form that does not resemble  $\sin(n\pi x)$ , we will need a more systematic approach to obtain the  $a_n$ . The standard way to do so is to invoke the "orthogonality" property of the eigenfunctions. For  $\{\sin(n\pi x)\}$ , we have

$$\begin{aligned} A_{m,n} &\equiv \int_0^1 \sin(n\pi x) \sin(m\pi x) dx = 0 \quad , \text{ if } m \neq n \\ &= 1/2 \quad , \text{ if } m = n \neq 0 \\ &= 1 \quad , \text{ if } m = n = 0 \end{aligned} \quad (1)$$

(Exercise: Verify that this is true.)

How to proceed:

**Step 1:** Multiply Eq. (14) in Slides #6 by  $\sin(m\pi x)$

$$\sin(m\pi x) \sum_{n=1}^{\infty} a_n \sin(n\pi x) = \sin(m\pi x) F(x) \quad (2)$$

**Step 2:** Integrate Eq. (2) over the whole domain, in our case from  $x = 0$  to  $x = 1$

$$\sum_{n=1}^{\infty} a_n \left( \int_0^1 \sin(m\pi x) \sin(n\pi x) dx \right) = \int_0^1 \sin(m\pi x) F(x) dx \quad ,$$

or,

$$\sum_{n=1}^{\infty} a_n A_{m,n} = \int_0^1 \sin(m\pi x) F(x) dx \quad , \quad (3)$$

where the value of  $A_{m,n}$  for given  $\{m, n\}$  can be obtained from Eq. (1).

The fact that  $A_{m,n}$  vanishes when  $m \neq n$  (i.e., two distinctive eigenfunctions are "orthogonal" to each other - we will explain this later) is critical for our scheme to work. This leads to our last step,

### Step 3:

For any given  $m$ , by noting that  $A_{m,n}$  vanishes when  $m \neq n$ , the infinite sum in the left hand side of Eq. (3) is reduced to a single term

$$\begin{aligned}\sum_{n=1}^{\infty} a_n A_{m,n} &= a_1 A_{m,1} + a_2 A_{m,2} + a_3 A_{m,3} + \dots + a_m A_{m,m} + \dots \\ &= 0 + 0 + 0 + \dots + a_m A_{m,m} + 0 + \dots \\ &= a_m A_{m,m} \\ &= a_m/2 \quad (\text{since } A_{m,m} = 1/2, \text{ from Eq. (1)})\end{aligned}\tag{4}$$

Using (3) and (4), we obtain a useful expression for the coefficients  $a_m$ ,

$$a_m = 2 \int_0^1 \sin(m\pi x) F(x) dx .\tag{5}$$

We are done! This is all we need to evaluate the coefficients from the  $F(x)$  in the initial condition.

## Example 1

Let's get back to the example in Slides #6, where  $F(x) = 4\sin(3\pi x) + 7\sin(8\pi x)$ . Using Eq. (5), we have

$$\begin{aligned} a_m &= 2 \int_0^1 \sin(m\pi x) F(x) dx \\ &= 2 \int_0^1 \sin(m\pi x) (4\sin(3\pi x) + 7\sin(8\pi x)) dx \\ &= 2 (4 A_{m,3} + 7 A_{m,8}) \end{aligned}$$


So, we have  $a_3 = 4$ ,  $a_8 = 7$ , and  $a_n = 0$  for all other  $n$ .

## Example 2

Let's try a more complicated initial condition for the Heat equation in Slides #6:

$$(iii) \quad u(x, 0) = F(x) \quad , \quad \text{where} \quad \begin{aligned} F(x) &= x \quad , \quad 0 \leq x \leq 1/2 \quad , \\ &= 1-x \quad , \quad 1/2 \leq x \leq 1 \end{aligned}$$

(This is a triangular-shaped distribution. See diagram below.)



In this case, using Eq. (5), we have


$$a_m = 2 \left[ \int_0^{1/2} \sin(m\pi x) x dx + \int_{1/2}^1 \sin(m\pi x) (1-x) dx \right] ,$$

$$\Rightarrow a_m = \frac{4}{m^2 \pi^2} \quad , \quad m = 1, 5, 9, 13, \dots$$

$$a_m = \frac{-4}{m^2 \pi^2} \quad , \quad m = 3, 7, 11, 15, \dots$$

$$a_m = 0 \quad , \quad \text{when } m \text{ is even .}$$

Plugging the  $a_m$  back to the solution of the Heat equation in Slides #6, we obtain the complete solution under the new initial state. The following is a plot of the solution  $u(x,t)$  at selected values of  $t$ .



Initial state (black);  
Solution at  $t = 0.01$  (gray), 0.03 (red), and 0.1 (green)

The solution is truncated at  $n = 10$