

Magnetic catalysis and chiral shift in dense matter

Igor Shovkovy*

*E.V. Gorbar, V.A. Miransky, I.S., Phys. Rev. C 80 (2009), 032801(R)

+ work in progress

Strong and Electroweak Matter 2010
McGill University, Montreal, Canada

Dense relativistic matter

- ⊙ Dense relativistic matter is common inside compact stars

- Electrons in white dwarfs

$$T \ll m \approx \mu \quad (\text{i.e., } T \approx 1 \text{ keV} \ \& \ \mu \approx 1 \text{ MeV})$$

- Neutrons of nuclear matter

$$T \ll m \approx \mu \quad (\text{i.e., } T \approx 10 \text{ MeV} \ \& \ \mu \approx 1 \text{ GeV})$$

- Electrons inside stellar nuclear matter

$$m \approx T \ll \mu \quad (\text{i.e., } T \approx 10 \text{ MeV} \ \& \ \mu \approx 100 \text{ MeV})$$

- Dense quark matter in stellar cores (if formed)

Dense relativistic matter

- ⦿ Dense relativistic matter is common inside compact stars

- Electrons in white dwarfs

$$T \ll m \approx \mu \quad (\text{i.e., } T \lesssim 1 \text{ keV} \ \& \ \mu \approx 1 \text{ MeV})$$

- Neutrons of nuclear matter

$$T \ll m \approx \mu \quad (\text{i.e., } T \lesssim 10 \text{ MeV} \ \& \ \mu \approx 1 \text{ GeV})$$

- Electrons inside stellar nuclear matter

$$m \lesssim T \ll \mu \quad (\text{i.e., } T \lesssim 10 \text{ MeV} \ \& \ \mu \approx 100 \text{ MeV})$$

- Dense quark matter in stellar cores (if formed)

General idea

- Topological current in relativistic matter in a magnetic field (3+1 dimensions)

$$\langle \bar{\psi} \gamma^3 \gamma^5 \psi \rangle = \frac{eB}{2\pi^2} \mu \quad (\text{free theory!})$$

[Metlitski, Zhitnitsky, PRD **72**, 045011 (2005)]

- Should there be a dynamical “mass” Δ , associated with this condensate?

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_\Delta \quad \text{where} \quad \mathcal{L}_\Delta \simeq \Delta \bar{\psi} \gamma^3 \gamma^5 \psi$$

- Note: $\Delta=0$ is not protected by any symmetry


Lesson from graphene

- Dynamics of Quantum Hall Effect in graphene (\approx QED in 2+1 dimensions)
 - Parity and time-reversal odd Dirac mass

$$\Delta \sim \langle \bar{\Psi} \gamma^3 \gamma^5 \Psi \rangle$$

[Gorbar, Gusynin, Miransky, I.S., PRB 78, 085437 (2008)]

- Δ describes the 0th plateau in Quantum Hall effect in graphene



Zhang et al., PRL 96, 136806 (2006)

Model

- Lagrangian density:

$$\mathcal{L} = \bar{\psi} (iD_\nu + \mu_0 \delta_\nu^0) \gamma^\nu \psi + \frac{G_{\text{int}}}{2} \left[(\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma^5\psi)^2 \right]$$

- The dimensionless coupling is

$$g \equiv G_{\text{int}} \Lambda^2 / (4\pi^2) \ll 1$$

- Magnetic field is inside


where

$$D_\nu = \partial_\nu - ieA_\nu \quad (\text{Landau gauge})$$

$$A_\nu = xB\delta_\nu^2$$

Approximation

- Gap equation in mean-field approximation:



$$G^{-1}(u, u') = S^{-1}(u, u') - iG_{\text{int}} \left\{ G(u, u) - \gamma^5 G(u, u) \gamma^5 - \text{tr}[G(u, u)] + \gamma^5 \text{tr}[\gamma^5 G(u, u)] \right\} \delta^4(u - u')$$

where

$$iG^{-1}(u, u') = \left[(i\partial_t + \mu)\gamma^0 - (\boldsymbol{\pi} \cdot \boldsymbol{\gamma}) - \pi^3 \gamma^3 + i\tilde{\mu}\gamma^1\gamma^2 + \Delta\gamma^3\gamma^5 - m \right] \delta^4(u - u')$$

and

$$iS^{-1}(u, u') = \left[(i\partial_t + \mu_0)\gamma^0 - (\boldsymbol{\pi} \cdot \boldsymbol{\gamma}) - \pi^3 \gamma^3 \right] \delta^4(u - u')$$

Vacuum state

- ⦿ Magnetic catalysis (spontaneous generation of a nonzero Dirac mass even at $g \ll 1$):

$$m_0^2 = \frac{1}{\pi l^2} \exp\left(-\frac{\Lambda^2 l^2}{g}\right) \text{ where } l = 1/\sqrt{|eB|}$$

(along with $\mu = \mu_0$)

[Gusynin, Miransky, I.S., PRL **73**, 3499 (1994); PLB **349**, 477 (1995)]

- ⦿ The solution exists for $\mu_0 < m_0$, although it will be less stable than the normal state ($m = 0$) already for

$$\mu_0 \gtrsim m_0/\sqrt{2} \text{ [Clogston, PRL } \mathbf{29}, 266 \text{ (1962)]}$$

“Abnormal” normal ground state

- ⊙ The gap equation allows another solution,

$$\mu \simeq \mu_0 \text{ and } \Delta \simeq g\mu_0 eB/\Lambda^2$$

- ⊙ This solution is almost independent of temperature when $T \ll \mu$
- ⊙ This is the normal ground state since its symmetry is same as in the Lagrangian
- ⊙ Besides, there is no trivial solution $\Delta=0$

Change of ground state

- The free energy in the state with $m \neq 0$ (broken symmetry)

$$\Omega_m \simeq -\frac{m_0^2}{2(2\pi l)^2} \left(1 + (m_0 l)^2 \ln |\Lambda l|\right)$$

- The free energy in the normal state, $\Delta = 0$

$$\Omega_\Delta \simeq -\frac{\mu_0^2}{(2\pi l)^2} \left(1 - g \frac{|eB|}{\Lambda^2}\right)$$

- So, indeed symmetry is restored for $\mu > \mu_c$,

$$\mu_c \simeq m_0 / \sqrt{2}$$


Physical meaning of Δ

- The dispersion relation of quasiparticles:


$$\omega_{n,\sigma} = -\mu \pm \sqrt{[k_3 + \sigma\Delta]^2 + 2n|eB|}$$

where $\sigma = \pm 1$ is the chirality





- Longitudinal momenta of opposite chirality fermions are *shifted*, i.e., $k_3 = k_3 \pm \Delta$
- All Landau levels ($n = 0$) are affected by Δ





Magnetic catalysis at $\mathbb{W}_0=0$



T=0 results



T ≠ 0 results



- These are smoothed versions of the $T=0$ results
- The dependence $\left[\frac{W}{M} \right] - \left[\frac{W}{M} \right]_0$ versus $\left[\frac{W}{M} \right]_0$ (not shown) at $T \neq 0$ is similar to Δ versus $\left[\frac{W}{M} \right]_0$ (shown)

Induced axial current

- ⦿ The axial current in the ground state is

$$\langle \bar{\psi} \gamma^3 \gamma^5 \psi \rangle = \frac{eB}{2\pi^2} \mu - \frac{|eB|}{2\pi^2} \Delta - \frac{|eB|}{\pi^2} \Delta \sum_{n=1}^{\infty} \kappa(\sqrt{2n|eB|}, \Lambda)$$

- ⦿ In addition to the topological contribution, $\frac{eB}{2\pi^2} \mu$ there are dynamical ones $\frac{|eB|}{2\pi^2} \Delta$
- ⦿ An equivalent result is also obtained in the Pauli-Villars regularization
- ⦿ **Note:** on the solution to the gap equation:

$$\langle j_5^3(u) \rangle = \frac{2\Delta}{G_{\text{int}}} = \frac{\Delta}{2\pi^2} \frac{\Lambda^2}{g}$$

Potential implications

- ⦿ Physical properties to be affected
 - transport
 - emission

(must be sensitive to anisotropy and/or CP violation)
- ⦿ Specific physical systems
 - Compact stars
 - Quark stars (quarks)
 - Hybrid stars (quarks, electrons)
 - Neutron stars (electrons)
 - White dwarfs (electrons)
 - Heavy ion collisions [Kharzeev & Zhitnitsky, NPA 797, 67(2007), Kharzeev, McLerran & Warringa, NPA 803, 227 (2008), ...]

Pulsar kicks

- The dynamical chiral shift parameter is driven by chemical potential (T μ)

$$\Delta \simeq g\mu_0 e B / \Lambda^2$$

and is almost independent of temperature

- This creates an anisotropy in the distribution of left-handed quarks/electrons
- The anisotropy is transferred to left-handed neutrinos by elastic scattering
- Pulsar gets a kick when neutrinos escape



Supernova explosions

- ⦿ Because of the robustness of the axial currents at finite temperatures, even supernova explosions may be affected
- ⦿ A small early-time neutrino asymmetry may *facilitate* explosions and give a kick at the same time, e.g., see
[Fryer & Kusenko, *Astrophys. J. Supp.* **163**, 335 (2006)]

Summary

- ⊙ $\mu < \mu_c$: Chiral symmetry is broken in the ground state (magnetic catalysis)
- ⊙ $\mu > \mu_c$: Normal ground state of dense relativistic matter in a magnetic field is characterized by
 - Chiral shift parameter (may have dramatic implications for stars)
 - Axial current along the field (physical effects are not obvious)
 - No solution with vanishing Δ exists

Outlook

- ⦿ Calculation of neutrino emission/diffusion in the state with a chiral shift parameter (work in progress)
- ⦿ Transport properties of the normal state with nonzero chiral shift parameter
- ⦿ The fate of the induced axial current in the renormalized models (work in progress)
- ⦿ Modification of the chiral magnetic effect due to “vector-like” Δ in heavy ion collisions

[Fukushima, Kharzeev & Warringa, PRD **78**, 074033 (2008)]

Thank you