

# TOPIC VII

## UNCONSTRAINED OPTIMIZATION II

### [1] Relative Maximum and Minimum

(1) Review for  $y = f(x)$ :

- FOC:  $f'(x) = 0 \Rightarrow$  get  $x^*$ .
- SOC:  $f''(x^*) > 0 \Rightarrow x^*$  is relative min. point.  
 $f''(x^*) < 0 \Rightarrow x^*$  is relative max. point.

(2) For  $y = f(x_1, x_2, \dots, x_n)$ :

- FOC:  $\frac{\partial f}{\partial x_1} \equiv f_1 = 0; \frac{\partial f}{\partial x_2} \equiv f_2 = 0; \dots; \frac{\partial f}{\partial x_n} \equiv f_n = 0.$

• SOC:

- From  $f_1 = \frac{\partial f}{\partial x_1}$ :

- $\frac{\partial^2 f}{\partial x_1 \partial x_1} = \frac{\partial}{\partial x_1} \left( \frac{\partial f}{\partial x_1} \right) = f_{11};$

- $\frac{\partial^2 f}{\partial x_1 \partial x_2} = \frac{\partial}{\partial x_2} \left( \frac{\partial f}{\partial x_1} \right) = f_{12};$

:

- $\frac{\partial^2 f}{\partial x_1 \partial x_n} = \frac{\partial}{\partial x_n} \left( \frac{\partial f}{\partial x_1} \right) = f_{1n};$

- From  $f_2 = \frac{\partial f}{\partial x_2}$ :

- $\frac{\partial^2 f}{\partial x_2 \partial x_1} = \frac{\partial}{\partial x_1} \left( \frac{\partial f}{\partial x_2} \right) = f_{21}$

- $\frac{\partial^2 f}{\partial x_2 \partial x_2} = \frac{\partial}{\partial x_2} \left( \frac{\partial f}{\partial x_2} \right) = f_{22}$

:

- $H_n = \begin{pmatrix} f_{11} & f_{12} & f_{13} & \cdots & f_{1n} \\ f_{21} & f_{22} & f_{23} & \cdots & f_{2n} \\ \vdots & \vdots & \vdots & & \\ f_{n1} & f_{n2} & f_{n3} & \cdots & f_{nn} \end{pmatrix}$  : Hessian matrix.

(3) Young's Theorem:

- $\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i} \quad (f_{ij} = f_{ji})$

## Digression on Matrix Theory:

Definition 1:

$A_{n \times n}$  is called symmetric, iff  $A^t = A$ .

- $[a_{ji}] = [a_{ij}] \Rightarrow a_{ji} = a_{ij}$

EX 1:  $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \\ 3 & 4 & 1 \end{pmatrix}$

EX 2:  $H_n$

- $f_{ij} = f_{ji} \Rightarrow H_n^t = H_n$ .

Definition 2: Quadratic form

- $x : n \times 1$  vector =  $\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$ ;  $A_{n \times n}$ : symmetric.

- Then,  $x'Ax$  is of a quadratic form in  $n$  variables,  $x_1, \dots, x_n$ .
- $\underset{1 \times n}{X^t} \underset{n \times n}{A} \underset{n \times 1}{X}$  is a scalar.

$$\text{EX 1: } \mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}; \mathbf{X} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}; \mathbf{A}: \text{symmetric}; a_{ij} = a_{ji}$$

$$\begin{aligned} \bullet \quad \mathbf{x}'\mathbf{A}\mathbf{x} &= a_{11}x_1^2 + 2a_{12}x_1x_2 + 2a_{13}x_1x_3 \\ &\quad + a_{22}x_2^2 + 2a_{23}x_2x_3 \\ &\quad + a_{33}x_3^2. \end{aligned}$$

$$\text{EX 2: } \mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \quad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, a_{12} = a_{21}$$

$$\bullet \quad \mathbf{x}'\mathbf{A}\mathbf{x} = a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2$$

$$\text{EX 3: } \mathbf{A} = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$$

$$\begin{aligned} \bullet \quad \mathbf{x}'\mathbf{A}\mathbf{x} &= 2x_1^2 + 2x_1x_2 + 2x_2^2 = 2(x_1^2 + x_1x_2) + 2x_2^2 \\ &= 2\left(x_1^2 + x_1x_2 + \frac{x_2^2}{4} - \frac{x_2^2}{4}\right) + 2x_2^2 \\ &= 2\left(x_1 + \frac{x_2}{2}\right)^2 + \frac{3}{2}x_2^2 > 0, \text{ unless } x_1 = x_2 = 0. \text{ (i.e., } \mathbf{x} = \mathbf{0}) \end{aligned}$$

EX 4:  $A = \begin{pmatrix} -2 & -1 \\ -1 & -2 \end{pmatrix}$

$$x'Ax = -2x_1^2 - 2x_1x_2 - 2x_2^2 = -2\left(x_1 + \frac{x_2}{2}\right)^2 - \frac{3}{2}x_2^2 < 0,$$

unless  $x = 0$

Definition 3:

- $A_{n \times n}$  is symmetric.
- If, for any  $x \neq 0$  (vector),  $x'Ax > 0$ ,  $A$  is called positive definite  
[If, for any  $x \neq 0$  (vector),  $x'Ax \geq 0$ ,  $A$  is called positive semidefinite.]
- If, for any  $x \neq 0$  (vector),  $x'Ax < 0$ ,  $A$  is called negative definite.  
[If, for any  $x \neq 0$  (vector),  $x'Ax \leq 0$ ,  $A$  is called negative semidefinite.]

EX:  $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$  p.d.;  $\begin{pmatrix} -2 & -1 \\ -1 & -2 \end{pmatrix}$  n.d.

## Definition 4: Principal Minors

- $A_{n \times n}$ .

- $|A_1| = a_{11}; |A_2| = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix}; |A_3| = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}; |A_4| = \begin{vmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{vmatrix}$ .

EX 1:  $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$

- $|A_1| = 2 > 0; |A_2| = \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} = 4 - 1 = 3 > 0$ .

EX 2:  $A = \begin{pmatrix} -2 & -1 \\ -1 & -2 \end{pmatrix}$

- $|A_1| = -2 < 0; |A_2| = (-2) \times (-2) - (-1) \times (-1) = 3 > 0$

**End of Digression**

### **Theorem:**

- $A_{n \times n}$  : symmetric.
- A is negative definite iff  $|A_1| < 0$  ;  $|A_2| > 0$  ;  $|A_3| < 0$ ;  $|A_4| > 0$  ; . . . .
- A: positive definite iff  $|A_1| > 0$  ;  $|A_2| > 0$  ;  $|A_3| > 0$ ; . . . .

### **Return to Hessian matrix:**

### **SOC:**

- For max,  $H_n$  should be negative definite.
- For min,  $H_n$  should be positive definite.


### **Sum-up:**

- FOC:  $f_1 = f_2 = f_3 = \dots = f_n = 0$
- SOC:
  - For max,  $|H_1| < 0$  ,  $|H_2| > 0$  ,  $|H_3| < 0$  , . . . .
  - For min,  $|H_1| > 0$  ,  $|H_2| > 0$  ,  $|H_3| > 0$  , . . . .

### **<Intuitive Proof>**

- $y = f(x_1, x_2)$ .
- God's world (we don't know):

$$x_1 = x_1(s); x_2 = x_2(s); y = f(x_1(s), x_2(s))$$



- FOC:

$$\frac{dy}{ds} = \frac{\partial f}{\partial x_1} \cdot \frac{dx_1}{ds} + \frac{\partial f}{\partial x_2} \cdot \frac{dx_2}{ds} \equiv f_1 \frac{dx_1}{ds} + f_2 \frac{dx_2}{ds} = 0 .$$

- Choose  $x_1, x_2$ , such that  $f_1 = f_2 = 0$ .
- Then, FOC holds, regardless of  $\frac{dx_1}{ds}, \frac{dx_2}{ds}$ .

- SOC:

$$\bullet \frac{dy}{ds} = f_1 \frac{dx_1}{ds} + f_2 \frac{dx_2}{ds} = f_1(x_1, x_2) \frac{dx_1}{ds} + f_2(x_1, x_2) \frac{dx_2}{ds} .$$

$$\bullet \frac{d^2y}{ds^2} = f_{11} \left( \frac{dx_1}{ds} \right)^2 + 2f_{12} \left( \frac{dx_1}{ds} \right) \left( \frac{dx_2}{ds} \right) + f_{22} \left( \frac{dx_2}{ds} \right)^2 .$$

$$= \begin{pmatrix} \frac{dx_1}{ds} & \frac{dx_2}{ds} \end{pmatrix} \begin{pmatrix} f_{11} & f_{12} \\ f_{21} & f_{22} \end{pmatrix} \begin{pmatrix} \frac{dx_1}{ds} \\ \frac{dx_2}{ds} \end{pmatrix} .$$



- For max  $y$ ,  $\frac{d^2y}{ds^2} < 0$ .

It happens regardless of  $\frac{dx_1}{ds}$ ,  $\frac{dx_2}{ds}$ , if  $H_2$  is negative definite.

- For min  $y$ ,  $\frac{d^2y}{ds^2} > 0$ .

It happens regardless of  $\frac{dx_1}{ds}$ ,  $\frac{dx_2}{ds}$ , if  $H_2$  is positive definite.

EX 1:

- $z = 8x^3 + 2xy - 3x^2 + y^2 + 1$

- FOC: •  $\frac{\partial z}{\partial x} = 24x^2 + 2y - 6x = 0$ .      ①

- $\frac{\partial z}{\partial y} = 2x + 2y = 0$ .      ②

- From ②:

$$y = -x \quad \text{③}$$

- ③ → ①;

$$24x^2 - 2x - 6x = 0$$

$$24x^2 - 8x = 0$$

$$3x^2 - x = 0$$

$$(3x - 1)x = 0$$

- $(x^*, y^*) = (0, 0)$  or  $(x^*, y^*) = (1/3, -1/3)$ .

- SOC:

- $\frac{\partial^2 z}{\partial x \partial x} = 48x - 6$ ;      •  $\frac{\partial^2 z}{\partial x \partial y} = 2$

- $\frac{\partial^2 z}{\partial y \partial x} = 2$ ;      •  $\frac{\partial^2 z}{\partial y \partial y} = 2$

- $H_2 = \begin{pmatrix} 48x - 6 & 2 \\ 2 & 2 \end{pmatrix}$

- When  $(x, y) = (1/3, -1/3)$ ,  $H_2 = \begin{pmatrix} 10 & 2 \\ 2 & 2 \end{pmatrix}$ .

- $|H_1| = 10 > 0$ ;  $|H_2| = 20 - 4 = 16 > 0$ .

- min. point:  $(x, y, z) = (1/3, -1/3, 28/29)$

- When  $(x, y) = (0, 0)$ ,  $H_2 = \begin{pmatrix} -6 & 2 \\ 2 & 2 \end{pmatrix}$ .

- $|H_1| = -6 < 0$ ;  $|H_2| = -12 - 4 = -16 < 0$

- Who knows? (saddle point)

EX 2:

- $z = x + 2ey - e^x - e^{2y}$

- FOC:

- $\frac{\partial z}{\partial x} = 1 - e^x = 0$

- $\frac{\partial z}{\partial y} = 2e - 2e^{2y} = 0$

- $(x^*, y^*, z^*) = (0, 1/2, -1)$ .

(Why?  $e - e^{2y} = 0 \rightarrow e = e^{2y} \rightarrow \ln e = \ln e^{2y} \rightarrow 1 = 2y$ )

- SOC:

- $\frac{\partial^2 z}{\partial x \partial x} = -e^x$ ;      •  $\frac{\partial^2 z}{\partial x \partial y} = 0$

- $\frac{\partial^2 z}{\partial y \partial y} = -4e^{2y}$ .

- $H_2 = \begin{pmatrix} -e^x & 0 \\ 0 & -4e^{2y} \end{pmatrix}$

- When  $x = 0$  and  $y = 1/2$ ,  $H_2 = \begin{pmatrix} -1 & 0 \\ 0 & -4e \end{pmatrix}$ .

- $|H_1| = -1 < 0$ ;  $|H_2| = 4e > 0 \Rightarrow$  local max.  $(x, y, z) = (0, 1/2, -1)$ .

EX 3:

- $z = 2x_1^2 + x_1x_2 + 4x_2^2 + x_1x_2 + x_1^2 + 2.$

- FOC:

- $\partial z/\partial x_1 = 4x_1 + x_2 + x_3 = 0$

- $\partial z/\partial x_2 = x_1 + 8x_2 = 0$

- $\partial z/\partial x_3 = x_1 + 2x_3 = 0$

$$\Rightarrow \begin{bmatrix} 4 & 1 & 1 \\ 1 & 8 & 0 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

- $x_1^* = x_2^* = x_3^* = 0.$

- SOC:

- $H_n = \begin{bmatrix} 4 & 1 & 1 \\ 1 & 8 & 0 \\ 1 & 0 & 2 \end{bmatrix}.$

- $|H_1| = 4 > 0;$

- $|H_2| = \begin{vmatrix} 4 & 1 \\ 1 & 8 \end{vmatrix} = 32 - 1 = 31 > 0;$

$$|H_3| = \begin{vmatrix} 4 & 1 & 1 \\ 1 & 8 & 0 \\ 1 & 0 & 2 \end{vmatrix} = 4 \cdot 8 \cdot 2 - 1 \cdot 8 \cdot 1 - 1 \cdot 1 \cdot 2 = 54 > 0.$$

- local min. point:  $(x_1, x_2, x_3, z) = (0, 0, 0, 2)$ .

EX 4:

- $z = -x_1^3 + 3x_1x_3 + 2x_2 - x_2^2 - 3x_3^2$ .

- FOC:

- $\partial z / \partial x_1 = -3x_1^2 + 3x_3 = 0$  ①

- $\partial z / \partial x_2 = 2 - 2x_2 = 0$  ②

- $\partial z / \partial x_3 = 3x_1 - 6x_3 = 0$  ③

- From ②:

- $x_2^* = 1$  ④

- $2 \cdot \textcircled{1} + \textcircled{3}$ :

- $-6x_1^2 + 3x_1 = 0 \rightarrow -2x_1^2 + x_1 = 0 \rightarrow x_1(-2x_1 + 1) = 0$

- $x_1^* = 0$  or  $x_1^* = 1/2$  ⑤

- By ③:

- $x_3 = (1/2)x_1$  ⑥

- $x_1^* = 0 \rightarrow x_3^* = 0; x_1^* = 1/2 \rightarrow x_3^* = 1/4$ .

- $(x_1^*, x_2^*, x_3^*, z^*) = (0, 1, 0, 1)$  or  $(1/2, 1, 1/4, 17/16)$ .

SOC:

- $\mathbf{H}_3 = \begin{bmatrix} -6x_1 & 0 & 3 \\ 0 & -2 & 0 \\ 3 & 0 & -6 \end{bmatrix}.$

- When  $x_1 = 1$ :

- $|\mathbf{H}_1| = 0 (\geq 0); |\mathbf{H}_2| = \begin{vmatrix} 0 & 0 \\ 0 & -2 \end{vmatrix} = 0 (\geq 0); |\mathbf{H}_3| = \begin{vmatrix} 0 & 0 & 3 \\ 0 & -2 & 0 \\ 3 & 0 & -6 \end{vmatrix} = 18 > 0.$

- Can't determine local max. or min. (In fact, local minimum.)

- When  $x_1 = 1/2$ :

- $\mathbf{H}_3 = \begin{bmatrix} -3 & 0 & 3 \\ 0 & -2 & 0 \\ 3 & 0 & -6 \end{bmatrix}.$

- $|\mathbf{H}_1| = -3; |\mathbf{H}_2| = \begin{vmatrix} -3 & 0 \\ 0 & -2 \end{vmatrix} = 6; |\mathbf{H}_3| = \begin{vmatrix} -3 & 0 & 3 \\ 0 & -2 & 0 \\ 3 & 0 & -6 \end{vmatrix} = -36 + 18 = -18.$

- max. point:  $(x_1, x_2, x_3, z) = (1/2, 1, 1/4, 17/16).$

## [2] Concavity Vs. Convexity

(1) Case for  $y = f(x)$ :

Definition:

A function  $y = f(x)$  is called strictly concave iff, for all  $x_1 \neq x_2$  and  $0 < \theta < 1$ ,  
 $f(\theta x_1 + (1-\theta)x_2) > \theta f(x_1) + (1-\theta)f(x_2)$ .


A function  $y = f(x)$  is called strictly convex iff, for all  $x_1 \neq x_2$  and  $0 < \theta < 1$ ,  
 $f(\theta x_1 + (1-\theta)x_2) < \theta f(x_1) + (1-\theta)f(x_2)$ .

Note:


A function  $y = f(x)$  is called concave iff, for all  $x_1 \neq x_2$  and  $0 < \theta < 1$ ,  
 $f(\theta x_1 + (1-\theta)x_2) \geq \theta f(x_1) + (1-\theta)f(x_2)$ .

A function  $y = f(x)$  is called strictly convex iff, for all  $x_1 \neq x_2$  and  $0 < \theta < 1$ ,  
 $f(\theta x_1 + (1-\theta)x_2) \leq \theta f(x_1) + (1-\theta)f(x_2)$ .

[Strictly Concave Function]




[Strictly Convex Function]






[Concave but Not Strict Concave Function]



[Convex but Not Strict Function]



Theorem:

For all  $x$ , if  $f''(x) < (>) 0$ ,  $f(x)$  is strictly concave (convex).

For all  $x$ , if  $f''(x) \leq (\geq) 0$ ,  $f(x)$  is concave (convex).

(2) Case for  $y = f(x_1, x_2, x_3, \dots, x_n)$ :

Definition:

- Choose any two different points  $u = (u_1, u_2, \dots, u_n)$  and  $v = (v_1, v_2, \dots, v_n)$ .

- Let  $y_u = f(u_1, u_2, \dots, u_n)$ ;

$$y_v = f(v_1, v_2, \dots, v_n);$$
$$y_\theta = f(\theta u_1 + (1-\theta)v_1, \theta u_2 + (1-\theta)v_2, \dots, \theta u_n + (1-\theta)v_n),$$

for any  $0 < \theta < 1$ .

- The function  $f(x_1, x_2, \dots, x_n)$  is strictly concave iff  $\theta y_u + (1-\theta)y_v < y_\theta$ .

[ $f(x_1, \dots, x_n)$  is concave iff  $\theta y_u + (1-\theta)y_v \leq y_\theta$ .]

- The function  $f(x_1, x_2, \dots, x_n)$  is strictly convex iff  $\theta y_u + (1-\theta)y_v > y_\theta$ .

[ $f(x_1, \dots, x_n)$  is convex iff  $\theta y_u + (1-\theta)y_v \geq y_\theta$ .]

Theorem:

- $y = f(x_1, \dots, x_n)$ .

- The function  $f$  is strictly concave iff  $H_n$  is nd for all points  $(x_1, \dots, x_n)$ .

[The function  $f$  is concave iff  $H_n$  is nsd for all points  $(x_1, \dots, x_n)$ .]

- The function  $f$  is strictly convex iff  $H_n$  is pd for all points  $(x_1, \dots, x_n)$ .

[The function  $f$  is concave iff  $H_n$  is psd for all points  $(x_1, \dots, x_n)$ .]

EX 1:

- $y = 2x - x^2$ .
- $y' = 2 - 2x \Rightarrow y'' = -2 < 0$ .
- $y = 2x - x^2$  is strictly concave.

EX 2:

- $y = f(x_1, x_2) = x_1^2 + x_2^2$ .
- $f_1 = 2x_1$ ;  
 $f_2 = 2x_2$
- $f_{11} = 2$ ;      •  $f_{12} = 0$   
 $f_{21} = 0$ ;      •  $f_{22} = 2$
- $H_n = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$  for all  $x_1$  and  $x_2$ .

$$\Rightarrow |H_1| = 2 > 0 ; |H_2| = 4 > 0$$

$\Rightarrow f$  is strictly convex.

Theorem:

- $y = f(x)$ .
- Suppose that  $f$  is concave (convex). If  $f'(x_0) = 0$ , then,  $y_0 = f(x_0)$  is a global maximum (minimum).

Theorem:

- $y = f(x_1, x_2, \dots, x_n)$ .

- Suppose that  $f$  is concave (convex).
- Suppose that at  $(x_1^*, \dots, x_n^*)$ ,  
 $f_1 = 0; f_2 = 0; \dots; f_n = 0$ .
- Then,  $y^* = f(x_1^*, \dots, x_n^*) =$  is a global maximum (minimum).

EX 1:

- $y = f(x_1, x_2) = x_1^2 + x_1 x_2 + x_2^2$ .
- FOC:
  - $f_1 = 2x_1 + x_2 = 0$
  - $f_2 = x_1 + 2x_2 = 0$
  - $\Rightarrow \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \Rightarrow x_1^* = x_2^* = 0$ .
- SOC:
  - $f_{11} = 2;$       •  $f_{12} = 1$
  - $f_{21} = 1;$       •  $f_{22} = 2$
  - $\Rightarrow H_n = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$ .
  - $\Rightarrow |H_1| = 2 > 0; |H_2| = 3 > 0$ , for all  $x_1$  and  $x_2$ .
- $y^* = f(x_1^*, x_2^*) = 0$  is a global minimum.

### [3] Firm's Behavior

#### (1) Assumptions:

- Production function:  $Q = f(L,K)$  .

- $MP_L = \partial Q / \partial L = f_L > 0$  .

- $MP_K = \partial Q / \partial K = f_K > 0$  .

- $f(L,K)$  is strictly concave:

- Hessian =  $\begin{bmatrix} f_{LL} & f_{LK} \\ f_{KL} & f_{KK} \end{bmatrix}$ .

- $f_{LL} < 0$  (Decreasing marginal product of labor.)

- $f_{LL}f_{KK} - (f_{LK})^2 > 0$ .

$[f_{LL}f_{KK} - (f_{LK})^2 > 0 \rightarrow f_{LL}f_{KK} > (f_{LK})^2 > 0 \rightarrow f_{KK} < 0$ , since  $f_{LL} < 0$ . Thus, the law of decreasing marginal product also applies to K.]

- The firm is perfectly competitive in both product and input markets.
  - Means that input and output prices are exogenously given.

#### (2) Profit maximization:

- $TR = pQ = Pf(L,K)$ .

- $TC = wL + rK$ .

- $\Pi = TR - TC$

$$= pf(L,K) - wL - rK.$$

- FOC:
  - $\partial\Pi/\partial L = pf_L - w = 0$
  - $\partial\Pi/\partial K = pf_K - r = 0$ .
- SOC:
  - $H_2 = \begin{bmatrix} pf_{LL} & pf_{LK} \\ pf_{LK} & pf_{KK} \end{bmatrix}$ 
    - $|H_1| = pf_{LL} < 0$  ;
    - $|H_2| = p^2 f_{LL} f_{KK} - p^2 (f_{LK})^2 = p^2 [f_{LL} f_{KK} - (f_{LK})^2] < 0$  for all L and K.
    - SOC for global max. holds.

(3) Implication of FOC:

- $pf_L = w \Rightarrow pMP_L = w \Rightarrow$  Marginal Revenue Product of Labor = wage.
  - Marginal Revenue Product of Labor: \$ you can make by hiring an extra unit of labor.
  - wage: \$ you should pay to hire an extra unit of labor.
- $pf_K = r \Rightarrow pMP_K = r \Rightarrow$  MRP of Capital = price of capital.

(4) Comparative Statics.

- Let  $L^*$  and  $K^*$  be the  $\Pi$ -max. input levels.
- Consider FOC:
  - $pf_L(L^*, K^*) - w = 0$
  - $pf_K(L^*, K^*) - r = 0$

- If  $r$ ,  $w$  or  $p$  changes,  $L^*$  and  $K^*$  changes.
  - Endogenous variables:  $L^*$ ,  $K^*$ .
  - Exogenous variables:  $p$ ,  $w$ , and  $r$ .
- Wish to find comparative statics.

1) Effect of  $w$ :

$$\bullet \begin{bmatrix} pf_{LL} & pf_{LK} \\ pf_{LK} & pf_{KK} \end{bmatrix} \begin{bmatrix} \frac{\partial L^*}{\partial w} \\ \frac{\partial K^*}{\partial w} \end{bmatrix} = \begin{bmatrix} -(-1) \\ -0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}.$$

$$\bullet \frac{\partial L^*}{\partial w} = \frac{\begin{vmatrix} 1 & pf_{LK} \\ 0 & pf_{KK} \end{vmatrix}}{|H_2|} = \frac{pf_{KK}}{|H_2|} = \frac{-}{+} < 0.$$

$$\bullet \frac{\partial K^*}{\partial w} = \frac{\begin{vmatrix} pf_{LL} & 1 \\ pf_{LK} & 0 \end{vmatrix}}{|H_2|} = \frac{-pf_{LK}}{|H_2|} = \frac{?}{+}.$$

2) Effect of  $r$ :

$$\bullet \begin{bmatrix} pf_{LL} & pf_{LK} \\ pf_{LK} & pf_{KK} \end{bmatrix} \begin{bmatrix} \frac{\partial L^*}{\partial r} \\ \frac{\partial K^*}{\partial r} \end{bmatrix} = \begin{bmatrix} -0 \\ -(-1) \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$\bullet \frac{\partial L^*}{\partial r} = \frac{\begin{vmatrix} 0 & pf_{LK} \\ 1 & pf_{KK} \end{vmatrix}}{|H_2|} = \frac{-pf_{LK}}{|H_2|} = \frac{?}{+}.$$

$$\bullet \frac{\partial K^*}{\partial r} = \frac{\begin{vmatrix} pf_{LL} & 0 \\ pf_{LK} & 1 \end{vmatrix}}{|H_2|} = \frac{pf_{LL}}{|H_2|} = \frac{-}{+} < 0.$$

3) Effect of p:

$$\bullet \begin{bmatrix} pf_{LL} & pf_{LK} \\ pf_{LK} & pf_{KK} \end{bmatrix} \begin{bmatrix} \frac{\partial L^*}{\partial p} \\ \frac{\partial K^*}{\partial p} \end{bmatrix} = = \begin{bmatrix} -f_L \\ -f_K \end{bmatrix}.$$

$$\bullet \frac{\partial L^*}{\partial p} = \frac{\begin{vmatrix} -f_L & pf_{LK} \\ -f_K & pf_{KK} \end{vmatrix}}{|H_2|} = \frac{-pf_L f_{KK} + pf_K f_{LK}}{|H_2|} = \frac{?}{+}.$$



$$\bullet \frac{\partial K^*}{\partial p} = \frac{\begin{vmatrix} pf_{LL} & -f_L \\ pf_{LK} & -f_K \end{vmatrix}}{|H_2|} = \frac{-pf_K f_{LL} + pf_L f_{LK}}{|H_2|} = \frac{?}{+}.$$

• But at least one should be positive. (Why?)

• Let  $L^* = L(p, w, r)$ ;  $K^* = K(p, w, r)$ .

•  $Q^* = f(L^*, K^*)$ .

• By Chain rule,

$$\begin{aligned} \partial Q^*/\partial p &= f_L(\partial L^*/\partial p) + f_K(\partial K^*/\partial p) \\ &= \frac{f_L(-pf_L f_{KK} + pf_K f_{LK}) + f_K(-pf_K f_{LL} + pf_L f_{LK})}{|H_2|} \\ &= \frac{-pf_{KK} f_L^2 + 2pf_{LK} f_L f_K - pf_{LL} f_K^2}{|H_2|} \\ &= \frac{-p}{|H_2|} [f_{KK}(f_L)^2 - 2f_{LK}(f_L)(f_K) + f_{LL}(f_K)^2] \\ &= \frac{-p}{|H_2|} (f_K, -f_L) \begin{bmatrix} f_{LL} & f_{LK} \\ f_{LK} & f_{KK} \end{bmatrix} \begin{bmatrix} f_K \\ -f_L \end{bmatrix} > 0. \end{aligned}$$

•  $\partial Q^*/\partial p = f_L(\partial L^*/\partial p) + f_K(\partial K^*/\partial p) > 0$

⇒ Either  $\partial L^*/\partial p$  or  $\partial K^*/\partial p$  should be negative.

⇒ If both are positive,  $\partial Q^*/\partial p < 0$ , which is a contradiction.

EX 1:

- $Q = AL^{1/3}K^{1/3}$ .
- $\Pi = pAL^{1/3}K^{1/3} - wL - rK$ .
- FOC:
  - $\partial\Pi/\partial L = (1/3)pAL^{-2/3}K^{1/3} - w = 0$ ;
  - $\partial\Pi/\partial K = (1/3)pAL^{1/3}K^{-2/3} - r = 0$ .

$$\Rightarrow (1/3)pAL^{-2/3}K^{1/3} = w; \quad \textcircled{1}$$

$$(1/3)pAL^{1/3}K^{-2/3} = r. \quad \textcircled{2}$$

①÷②:

$$\frac{L^{-2/3}K^{1/3}}{L^{1/3}K^{-2/3}} = \frac{w}{r}$$

$$\Rightarrow L^{-1}K = w/r$$

$$\Rightarrow L = (w/r)L \quad \textcircled{3}$$

③→①:

$$(1/3)pAL^{-2/3}[(w/r)L]^{1/3} = w$$

$$\Rightarrow (1/3)pA(w/r)^{1/3}L^{-1/3} = w$$

$$\Rightarrow (1/3)pA(w/r)^{1/3} = wL^{1/3}$$

$$\Rightarrow (1/3)^3 p^3 A^3 (w/r) = w^3 L$$

$$\Rightarrow L^* = (1/27)p^3 A^3 / (w^2 r) \quad \textcircled{4}$$

④→③:

$$K^* = (w/r)L^* = (1/27)p^3 A^3 / (w^2 r) = (1/27)p^3 A^3 / (r^2 w). \quad \textcircled{5}$$

- SOC:

- $$H_2 = \begin{bmatrix} -\frac{2}{9}pAL^{-5/3}L^{1/3} & \frac{1}{9}pAL^{-2/3}K^{-2/3} \\ \frac{1}{9}pAL^{-2/3}K^{-2/3} & -\frac{2}{9}pAL^{1/3}K^{-5/2} \end{bmatrix}.$$

- $|H_1| = \frac{2}{9}pAL^{-5/3}K^{1/3} < 0$  for all L and K (> 0).

$$\begin{aligned} |H_2| &= \frac{4}{81}p^2A^2L^{-4/3}K^{-4/3} - \frac{1}{81}p^2A^2L^{-4/3}K^{-4/3} \\ &= \frac{3}{81}p^2A^2L^{-4/3}K^{-4/3} > 0 \text{ for all L and K (>0)} \end{aligned}$$

- $L^* = (1/27)p^3A^3/(w^2r)$  and  $K^* = (1/27)p^3A^3/(r^2w)$  are global  $\Pi$ -max. input levels.