

Point-reflections in metric planes

Victor Pambuccian

Department of Integrative Studies, Arizona State University West
Phoenix, AZ 85069-7100, U.S.A. e-mail: pamb@math.west.asu.edu

Abstract

We axiomatize the class of groups generated by the point-reflections of a metric plane with a non-Euclidean metric, the structure of which turns out to be very rich compared to the Euclidean metric case, and state an open problem.

Mathematics Subject Classification: 51M05, 51M10, 51F15, 03B30.

1 Introduction

There is a very large literature on characterizations of groups of motions in terms of line-reflections or hyperplane-reflections (see [1]), but relatively little about groups generated by point-reflections. This subject has received some attention much later, in [4], [5], [6] (and, in a different setting, with an added differential structure, in e. g. [2] or [3]).

The purpose of this paper is to determine the theories of point-reflections that one obtains from the groups of isometries of Bachmann's metric planes.

If the metric plane is elliptic, i. e. if there are three line-reflections whose product is the identity, then the point-reflections coincide with the line-reflections, so that the axiom system of the group generated by point-reflections is identical to the one expressed in terms of line-reflections. The interesting case is thus that of non-elliptic metric planes.

2 Non-elliptic metric planes

2.1 Axiom system in terms of line-reflections

We shall first present non-elliptic metric planes as they appear in [1]. Our language will be a one-sorted one, with variables to be interpreted as 'rigid motions', containing a unary predicate symbol G , with $G(x)$ to be interpreted as ' x is a line-reflection', a constant symbol 1 , to be interpreted as 'the identity', and a binary operation \circ , with $\circ(a, b)$, which we shall write as $a \circ b$, to be interpreted as 'the composition of a with b '.

To improve the readability of the axioms, we introduce the following abbreviations:

$$\begin{aligned}
a^2 & :\Leftrightarrow a \circ a, \\
\iota(g) & :\Leftrightarrow g \neq 1 \wedge g^2 = 1, \\
a|b & :\Leftrightarrow G(a) \wedge G(b) \wedge \iota(a \circ b), \\
J(abc) & :\Leftrightarrow \iota((a \circ b) \circ c), \\
pq|a & :\Leftrightarrow p|q \wedge G(a) \wedge J(pqa).
\end{aligned}$$

The axioms are (we omit universal quantifiers whenever the axioms are universal sentences):

$$\mathbf{B 1} \quad (a \circ b) \circ c = a \circ (b \circ c)$$

$$\mathbf{B 2} \quad (\forall a)(\exists b) b \circ a = 1$$

$$\mathbf{B 3} \quad 1 \circ a = a$$

$$\mathbf{B 4} \quad G(a) \rightarrow \iota(a)$$

$$\mathbf{B 5} \quad G(a) \wedge G(b) \rightarrow G(a \circ (b \circ a))$$

$$\mathbf{B 6} \quad (\forall abcd)(\exists g) a|b \wedge c|d \rightarrow G(g) \wedge J(abg) \wedge J(cdg)$$

$$\mathbf{B 7} \quad ab|g \wedge cd|g \wedge ab|h \wedge cd|h \rightarrow (g = h \vee a \circ b = c \circ d)$$

$$\mathbf{B 8} \quad \bigwedge_{i=1}^3 pq|a_i \rightarrow G(a_1 \circ (a_2 \circ a_3))$$

$$\mathbf{B 9} \quad \bigwedge_{i=1}^3 g|a_i \rightarrow G(a_1 \circ (a_2 \circ a_3))$$

$$\mathbf{B 10} \quad (\exists ghj) g|h \wedge G(j) \wedge \neg j|g \wedge \neg j|h \wedge \neg J(jgh)$$

$$\mathbf{B 11} \quad (\forall x)(\exists ghj) G(g) \wedge G(h) \wedge G(j) \wedge (x = g \circ h \vee x = g \circ (h \circ j))$$

$$\mathbf{B 12} \quad G(a) \wedge G(b) \wedge G(c) \rightarrow a \circ (b \circ c) \neq 1$$

Since $a \circ b$ with $a|b$ represents a point-reflection, we may think of an unordered pair (a, b) with $a|b$ as a *point*, an element a with $G(a)$ as a *line*, two lines a and b for which $a|b$ as a pair of perpendicular lines, and say that a point (p, q) is *incident* with the line a if $pq|a$. With these figures of speech in mind, the above axioms make the following statements: B1, B2, and B3 are the group axioms for the operation \circ ; B4 states that line-reflections are involutions; B5 states the invariance of the set of line-reflections, B6 states that any two points can be joined by a line, which is unique according to B7 (we shall denote the line joining the points (a, b) and (c, d) by $\langle\langle a, b \rangle, \langle c, d \rangle\rangle$); B8 and B9 state that the composition of three reflections in lines that have a common point or a common perpendicular is a line-reflection; B10 states that there are three lines g, h, j such that g and h are perpendicular, but j is perpendicular to neither g nor h , nor does it go through the intersection point of g and h ; B11 states that every motion is the composition of two or three line-reflections, and B12 states that the composition of three line-reflections is never the identity. The function of the last axiom, B12, is to exclude elliptic geometries, and thus to ensure that the perpendicular from a point not on a line to that line is unique. The theory of non-elliptic metric planes, axiomatized by $\{B1 - B12\}$ will be denoted by \mathcal{B} .

2.2 Axiom system in terms of ternary geometric operations

The same class of models can also be axiomatized in the following manner: the language \mathcal{L} contains only one sort of individual variables, to be interpreted as ‘points’, three individual constants a_0, a_1, a_2 , to be interpreted as three non-collinear points, and two operation symbols, F and π . $F(abc)$ is the foot of the perpendicular from c to the line ab , if $a \neq b$, and a itself if $a = b$, and $\pi(abc)$ is the fourth reflection point whenever a, b, c are collinear points with $a \neq b$ and $b \neq c$, and arbitrary otherwise. By ‘fourth reflection point’ we mean the following: if we designate by σ_x the mapping defined by $\sigma_x(y) = \sigma(xy)$, i. e. the reflection of y in the point x , then, if a, b, c are three collinear points, by [1, §3,9, Satz 24b], the composition (product) $\sigma_c\sigma_b\sigma_a$, is the reflection in a point, which lies on the same line as a, b, c . That point is designated by $\pi(abc)$.

In order to formulate the axioms in a more readable way, we shall use the following abbreviations:

$$\sigma(ab) := \pi(aba), \quad (1)$$

$$R(abc) := \sigma(F(abc)c), \quad (2)$$

$$L(abc) :\leftrightarrow F(abc) = c \vee a = b, \quad (3)$$

where σ has the same meaning as above, $R(abc)$ stands for the reflection of c in ab (a line if $a \neq b$, the point a if $a = b$), and $L(abc)$ stands for ‘the points a, b, c are collinear (but not necessarily distinct)’.

The axiom system consists of the following axioms

$$\mathbf{C\ 1} \quad F(aab) = a$$

$$\mathbf{C\ 2} \quad \sigma(aa) = a$$

$$\mathbf{C\ 3} \quad \sigma(a\sigma(ab)) = b$$

$$\mathbf{C\ 4} \quad L(aba)$$

$$\mathbf{C\ 5} \quad L(abc) \rightarrow L(cba) \wedge L(bac)$$

$$\mathbf{C\ 6} \quad L(ab\sigma(ab))$$

$$\mathbf{C\ 7} \quad L(abF(abc))$$

$$\mathbf{C\ 8} \quad \sigma(ax) = \sigma(bx) \rightarrow a = b$$

$$\mathbf{C\ 9} \quad a \neq b \wedge F(abx) = F(aby) \rightarrow L(xyF(abx))$$

$$\mathbf{C\ 10} \quad a \neq b \wedge c \neq d \wedge F(abc) = c \wedge F(abd) = d \rightarrow F(abx) = F(cdx)$$

$$\mathbf{C\ 11} \quad \neg L(abx) \wedge F(xF(abx)y) = y \rightarrow F(abx) = F(aby)$$

$$\mathbf{C\ 12} \quad a \neq b \wedge a \neq c \wedge F(abc) = a \rightarrow F(acb) = a$$

$$\mathbf{C\ 13} \quad a \neq x \wedge x \neq y \wedge F(axy) = x \rightarrow F(a\sigma(ax)\sigma(ay)) = \sigma(ax)$$

$$\mathbf{C\ 14} \quad \sigma(\sigma(xa)\sigma(xb)) = \sigma(x\sigma(ab))$$

$$\mathbf{C\ 15} \quad u \neq v \wedge a \neq b \wedge F(abc) = a \rightarrow F(R(uva)R(uvb)R(uvc)) = R(uva)$$

$$\mathbf{C\ 16} \quad \neg L(oab) \wedge \neg L(abc) \rightarrow \sigma(F(xR(ocR(obR(oax)))o)x) = R(ocR(obR(oax)))$$

$$\mathbf{C\ 17} \quad \neg L(oab) \wedge \neg L(abc) \wedge \sigma(mx) = R(ocR(obR(oax))) \\ \wedge \sigma(ny) = R(ocR(obR(oay))) \rightarrow L(omn)$$

$$\mathbf{C\ 18} \quad a \neq b \wedge b \neq c \wedge F(abc) = c \wedge a \neq a' \wedge b \neq b' \wedge c \neq c' \wedge F(aba') = a \wedge F(bab') \\ = b \wedge F(cbc') = c \rightarrow \sigma(F(xR(cc'R(bb'R(aa'x)))\pi(abc))x) \\ = R(cc'R(bb'R(aa'x))) \wedge F(\pi(abc)cF(xR(cc'R(bb'R(aa'x)))\pi(abc))) = \pi(abc)$$

$$\mathbf{C\ 19} \quad \neg L(a_0a_1a_2)$$

The axioms make the following statements: C1 defines the value of $F(abc)$ when $a = b$ — it is an axiom with no geometric function (we could have opted to leave it undefined, but that would have lengthened the statements of the axioms C16 and C18); C2: the point a is a fixed point of the reflection σ_a , C3: reflections in points are involutory transformations (or the identity); C8: reflections of a point in two different points do not coincide; C4: a lies on the line determined by a and b ; C5: collinearity of three points is a symmetric relation; C6: the reflection of b in a is collinear with a and b ; C7: for $a \neq b$, the foot of the perpendicular from c to the line ab lies on that line; C9 states the uniqueness of the perpendicular to the line ab in the point $F(abc)$; C10: the foot of the perpendicular from x to the line ab does not depend on the particular choice of points a and b that determine the line ab ; C11: if x is a point outside of the line ab , and y is a point on the perpendicular from x to ab , then the feet of the perpendiculars of x and y to the line ab coincide; C12 states that perpendicularity is a symmetric relation (if ca is perpendicular to ab , then ba is perpendicular to ac); C13: if yx is perpendicular to xa , then so are $\sigma_a(y)\sigma_a(x)$ and $\sigma_a(x)a$; C14: reflections in points preserve midpoints; C15: reflections in lines preserve the orthogonality relation; C16 and C17 together state the three reflections theorem for lines having a point in common; C18 is the three reflections theorem for lines having a common perpendicular; C19: a_0, a_1, a_2 are three non-collinear points. With $\Sigma = \{\text{C1-C19}\}$, we proved in [8] the following

Theorem 1 Σ is an axiom system for non-elliptic metric planes. In every model of Σ , the operations F and π have the intended interpretations.

3 Axiom system for metric planes with non-Euclidean metric in terms of point-reflections

We now turn to yet another axiomatization of non-elliptic metric planes with non-Euclidean metric (i.e. in which there exists no rectangle), in terms of motions which are products of point-reflections, the individual constant 1, and the binary operation \circ , with $a \circ b$ standing for the composition of the motions a and b . In case a is a point-reflection, we will refer to a as a ‘point’ as well.




Figure 1: The definition of perpendicularity in terms of σ

To improve the readability of the axioms we introduce the following abbreviations:

$$\begin{aligned}
P(a) & :\Leftrightarrow a \neq 1 \wedge a \circ a = 1 \\
P(a_1, \dots, a_n) & :\Leftrightarrow \bigwedge_{i=1}^n P(a_i) \\
L(abc) & :\Leftrightarrow (a \circ b) \circ c = (c \circ b) \circ a \\
\sigma(ab) & := (a \circ b) \circ a \\
\varphi(eabcd) & :\Leftrightarrow (\neg L(abc) \wedge L(abd) \wedge L(cde) \wedge \sigma(e\sigma(bc)) = \sigma(\sigma(db)c)) \\
& \quad \vee (a \neq b \wedge L(abc) \wedge d = c) \vee (a = b \wedge d = a) \\
\pi(abc) & := c \circ (b \circ a) \\
\rho(eabcd) & :\Leftrightarrow \varphi(eabcu) \wedge d = \sigma(uc).
\end{aligned}$$

Here $P(a)$ stands for ‘ a is a point-reflection’, given that, in the group generated by point-reflections the only involutory elements are the point-reflections themselves. The subsequent abbreviations will be used only when all the variables that appear in them are point-reflections. $L(abc)$ stands for ‘ a, b, c are collinear’; $\sigma(ab)$ is the point obtained by reflecting b in a ; $\varphi(eabcd)$ holds, in case $a \neq b$, if d is the foot of the perpendicular from c to ab (as shown, for a, b, c not collinear in [9, Prop. 1] (e is a point needed in this construction, see Fig. 1)) and, in case $a = b$, if $d = a$; and $\rho(eabcd)$ stands for ‘ d is the reflection of c in the line ab if $a \neq b$ or in point a if $a = b$ ’.

The axioms for this axiom system are: B1, B2, B3, axiom P1, which ensures that, whenever a, b, c are collinear points, $\pi(abc)$ is a point as well, the axioms P2 and P3 stating the existence and uniqueness of the foot of the perpendicular from point c to line ab , whenever c does not lie on the line ab , as well as P4-P14, which are slightly changed variants of C8-C19.

P 1 $P(a, b, c) \rightarrow a \circ b \neq c$

P 2 $(\forall abc)(\exists de) P(a, b, c) \wedge \neg L(abc) \rightarrow P(e, d) \wedge \varphi(eabcd)$

- P 3** $P(a, b, c, d, e, d', e') \wedge \neg L(abc) \wedge \varphi(eabcd) \wedge \varphi(e'abcd') \rightarrow d = d'$
- P 4** $P(a, b, x) \wedge \sigma(ax) = \sigma(bx) \rightarrow a = b$
- P 5** $P(a, b, d, e, f, x, y) \wedge a \neq b \wedge \varphi(eabxd) \wedge \varphi(fabyd) \rightarrow L(xyd)$
- P 6** $P(a, b, c, d, e, f, u, v, x) \wedge a \neq b \wedge c \neq d \wedge L(abc) \wedge L(abd) \wedge \varphi(eabxu) \wedge \varphi(fcdxv) \rightarrow u = v$
- P 7** $P(a, b, d, e, f, u, v, x) \wedge \neg L(abx) \wedge \varphi(eabxu) \wedge L(xyu) \wedge \varphi(fabyv) \rightarrow u = v$
- P 8** $P(a, b, c, x, y, u) \wedge \neg L(abc) \wedge \varphi(xabca) \wedge \varphi(yacbu) \rightarrow u = a$
- P 9** $P(a, e, f, u, x, y) \wedge \neg L(axy) \wedge \varphi(eaxyx) \wedge \varphi(fa\sigma(ax)\sigma(ay)u) \rightarrow u = \sigma(ax)$
- P 10** $P(a, b, c, a', b', c', e, m, n, p, q, u, v, x) \wedge \neg L(abc) \wedge \varphi(eabca) \wedge u \neq v \wedge \varrho(nuva a') \wedge \varrho(puvbb') \wedge \varrho(quvcc') \wedge \varphi(ma'b'c'x) \rightarrow x = a'$
- P 11** $P(o, a, b, c, m, n, p, q, x, y, z, u, v) \wedge \neg L(oab) \wedge \neg L(obc) \wedge \varrho(moaxy) \wedge \varrho(nobyz) \wedge \varrho(pocz u) \wedge \varphi(qxuov) \rightarrow \sigma(vx) = u$
- P 12** $P(o, a, b, c, m, n, p, m', n', p', x, y, z, u, x', y', z', u', t, t') \wedge \neg L(oab) \wedge \neg L(obc) \wedge \varrho(moaxy) \wedge \varrho(nobyz) \wedge \varrho(pocz u) \wedge \varrho(m'oa x'y') \wedge \varrho(n'oby'z') \wedge \varrho(p'ocz'u') \wedge \sigma(tx) = u \wedge \sigma(t'x') = u' \rightarrow L(ott')$
- P 13** $P(a, b, c, a', b', c', m, n, p, t, u, v, w, x, y, z, w, o, g) \wedge L(abc) \wedge \neg L(aba') \wedge \neg L(bab') \wedge \neg L(cbc') \wedge \varphi(maba'a) \wedge \varphi(nbab'b) \wedge \varphi(pcbc'c) \wedge \varrho(taa'xy) \wedge \varrho(ubb'yz) \wedge \varrho(vcc'zw) \wedge \varphi(qxw\pi(abc)o) \wedge \varphi(r\pi(abc)cog) \rightarrow \sigma(ox) = w \wedge g = \pi(abc)$
- P 14** $(\exists abc) P(a, b, c) \wedge \neg L(abc)$

Finally, we need an axiom that lets us know that every rigid motion is a product of point-reflections. We do not know whether every element of the subgroup generated by point-reflections of the motion group G of a metric plane can be written as a product of at most a certain fixed number of point-reflections (whereas we do know that every element of G can be written as a product of at most three line-reflections). Unless we establish that there is an upper bound on the number of point-reflections needed (or that such an upper bound does not exist), we cannot determine the first-order theory of the group generated by point-reflections (as it may be either (1) the theory axiomatized by the axioms {B1 – B3, P1 – P14} in case there are, for every natural number k , products of point-reflections that cannot be written as a product of at most k point-reflections, or (2) the theory axiomatized by those axioms and an axiom stating that every rigid motion is a product of at most k point-reflections, should k be the least number with this property).¹ What we can do is to determine the $L_{\omega_1\omega}$ -theory of point-reflections, i. e. to state that every rigid motion is a product of an unspecified number of point-reflections as an infinite disjunction of first-order formulas. The axiom thus is

¹It was shown in [11] that, under additional assumptions, it is possible to write every product of point-reflections as a product of at most 4 point-reflections, but no such reduction is known in the general non-Euclidean metric case.

P 15 $(\forall a) \bigvee_{n=1}^{\infty} (\exists p_1 \dots p_n) P(p_1, \dots, p_n) \wedge a = p_1 \circ (\dots \circ p_n) \dots)$

Let $\Pi = \{\text{B1} - \text{B3}, \text{P1} - \text{P15}\}$. Given that we can define F in terms of φ , given that P2 and P3 ensure the existence and uniqueness of the value of $F(abc)$ for c not on ab , that C1-C7, C14 follow from our definitions of L , F and σ , and that the axioms C8-C19 follow from their translations P4-P14 into our language, we deduce that in every model of Π the individual variables x for which $P(x)$ holds can be interpreted as points, the defined notions F and π have the desired geometric interpretation, and thus, that the resulting structure is a non-elliptic metric plane with non-Euclidean metric (the metric cannot be Euclidean, given that, by P14 there are three non-collinear points a, b, c , and by P2, P3, there is precisely one point d on the line ab for which a point e with $\varphi(eabcd)$ exists; had the metric been Euclidean, then all points would have been “collinear” and for every point d on ab there would have been a point e with $\varphi(eabcd)$).

Let \mathfrak{M} be a model of Π . We now associate to every element $a \in \mathfrak{M}$ a mapping of the set S of points, i. e. of those members x of the universe of \mathfrak{M} for which $P(x)$ holds, into itself, which we denote by \tilde{a} and define by $\tilde{a}(x) := a \circ x \circ a^{-1}$. Note that $\widetilde{a \circ b} = \tilde{a} \cdot \tilde{b}$, where by \cdot we have denoted the operation of composition of maps. If $a \in S$, then $\tilde{a}(x) = \sigma(ax)$, so the set $\{\tilde{a} : P(a)\}$ generates a group \mathfrak{G} inside $Sym(S)$, which is precisely the group generated by the point-reflections of a non-elliptic metric plane with non-Euclidean metric (given that we know that σ has the desired interpretation). Given P15, the map $\tilde{}$ defines an isomorphism of \mathfrak{M} onto \mathfrak{G} . We have shown that

Theorem 2 Π is an $L_{\omega_1\omega}$ -axiom system for the group generated by the point-reflections of non-elliptic metric planes with non-Euclidean metric.

We now turn to metric planes whose metric is Euclidean, also called ‘metric-Euclidean planes’.

4 Point-reflections in metric-Euclidean planes

If axiom R (“There exists a rectangle” (see [1, §6,7])) holds in a metric plane, i. e.

$$(\exists abcd) a \neq b \wedge c \neq d \wedge a|c \wedge a|d \wedge b|c \wedge b|d,$$

is added to \mathcal{B} , then the group generated by point-reflections can be described very simply by means of B1-B3 and

E 1 $(\exists ab) P(a, b) \wedge a \neq b$

E 2 $P(a, b) \wedge a \circ b = b \circ a \rightarrow a = b$

E 3 $(\forall x)(\exists ab) P(a, b) \wedge (x = a \vee x = a \circ b)$

E 4 $P(a, b, c) \rightarrow P(a \circ (b \circ c))$

Let \mathfrak{M} be a model of B1-B3, E1-E4, let M be its universe, and let $P = \{m \in M : P(m)\}$ be the set of points in M . We define on $P \times P$ an equivalence relation \sim by $(a, b) \sim (c, d)$ if and only if $a \circ c = b \circ d$, and denote by $[a, b]$ the equivalence class of (a, b) . Let $G := P \times P / \sim$. We define on G an addition operation $+$ by $[a, b] + [c, d] := [a, b \circ c \circ d]$, which turns G into an Abelian group, as can be easily checked. We fix a point o in P , and consider all elements in G written as $[o, x]$ (notice that $[a, b] = [o, b \circ (a \circ o)]$). Writing \mathbf{x} for $[o, x]$, we check that $\sigma(\mathbf{a}\mathbf{b}) = 2\mathbf{a} - \mathbf{b}$ (i. e. that $[o, \sigma(ab)] + [o, b] = [o, a] + [o, a]$). By E2 we know that G must satisfy $2x = 0 \rightarrow x = 0$.

Any metric-Euclidean plane can be embedded in a Gaussian plane associated with the pair of fields (K, L) , where $K \subset L$, $[L : K] = 2$ (a generalization of the Gauss plane over (\mathbb{C}, \mathbb{R}) , see [7]), the points being elements of L , and thus the algebraic representation of point-reflections $\tilde{\mathbf{x}}$ is given by $\tilde{\mathbf{x}}(\mathbf{y}) = 2\mathbf{x} - \mathbf{y}$ and $(\tilde{\mathbf{x}} \cdot \tilde{\mathbf{y}})(\mathbf{z}) = 2(\mathbf{x} - \mathbf{y}) + \mathbf{z}$. Given that the only operations involved in the description of point-reflections and their composition is $+$ and $-$, every first-order sentence that is true in all metric-Euclidean planes must hold over arbitrary Abelian groups which satisfy $2x = 0 \rightarrow x = 0$ as well. Thus

Theorem 3 *{B1 – B3, E1 – E4} is an axiom system for the group generated by the point-reflections of metric-Euclidean planes.*

Related results have been proved in [10].

Acknowledgement. This paper was written while enjoying the hospitality of the Institute of Mathematics of the University of Erlangen, as a guest of Prof. Dr. Karl Strambach.

References

- [1] F. Bachmann, Aufbau der Geometrie aus dem Spiegelungsbegriff, 2. Auflage, Springer-Verlag, Berlin 1973.
- [2] Á. Figula, 3-dimensional Bol loops as sections in non-solvable Lie groups, Forum Math. 17 (2005), 431–460.
- [3] Á. Figula, K. Strambach, Affine extensions of loops, Abh. Math. Sem. Univ. Hamburg 74 (2004), 151–162
- [4] E. Gabrieli, H. Karzel, Point reflection geometries, geometric K-loops and unitary geometries, Resultate Math. 32 (1997), 66–72.
- [5] H. Karzel, A. Konrad, Reflection groups and K-loops, J. Geom. 52 (1995), 120–129.
- [6] H. Karzel, H. Wefelscheid, A geometric construction of the K-loop of a hyperbolic space, Geom. Dedicata 58 (1995), 227–236.
- [7] V. Pambuccian, Fragments of Euclidean and hyperbolic geometry, Sci. Math. Jpn. 53 (2001), 361–400.
- [8] V. Pambuccian, Constructive axiomatization of non-elliptic metric planes, Bull. Polish Acad. Sci. Math. 51 (2003), 49–57.
- [9] V. Pambuccian, Hyperbolic geometry in terms of point-reflections or of line-orthogonality, Math. Pannon. 15 (2004), 241–258.

- [10] W. Schwabhäuser, Modellvollständigkeit der Mittelpunktsgeometrie und der Theorie der Vektorgruppen, *Abh. Math. Sem. Univ. Hamburg* **48** (1979), 213–224.
- [11] C. Stroscher, Reduktion von Punktspiegelungsprodukten in der ebenen metrischen Geometrie, *Arch. Math.* **33** (1979), 183–192.