

The dynamics of N and P during *Ambrosia* litter degradation in urban versus rural environments

M. Christman¹, S. Hall², and B. Ball³


¹School of Sustainability, Arizona State University, PO Box 875502, Tempe, AZ 85287-4502, ²School of Life Sciences, Arizona State University, PO Box 874501, Tempe, AZ 85287-4501, and ³School of Mathematics and Natural Sciences, Arizona State University West Campus, PO Box 37100, Phoenix, AZ 85069-7100

Introduction:

-In fast urbanizing arid ecosystems, nitrogen (N) deposition is increasing due to anthropogenic sources.

-Previous studies show that N deposition is significantly higher in the Phoenix city core than in surrounding areas and that processes that promote N retention in other ecosystems are qualitatively different or absent in deserts, and that multiple N-loss pathways are prevalent.

-Excess N taken up by plants gets recycled through decomposition, but we don't know much about nutrient dynamics during photodegradation



Methodology:

-A litter decomposition experiment was conducted to explore the possible fates of added N temporarily stored in herbaceous vegetation.

-Over 40 weeks of decomposition, nitrogen, carbon, phosphorous, lignin, and microbial biomass were observed to represent both the chemical and biological dynamics during litter degradation.

-*Ambrosia deltoidea* litter was collected from city or downwind sites from plots receiving N+P treatments or plots receiving no nutrient addition.

-3 g of litter from each treatment and site was then placed into either UV transparent or UV opaque litter bags.

-Bags were then placed in the field at the Salt River Recreation area and the South Mountain East site, with litter from the city core or downwind, respectively.


-Samples were collected at intervals of 10, 20, 30, and 40 weeks and tested to determine whether differences in nutrient release during biological and photodegradation differs between the urban and rural ecosystem.



Results:

-Mass loss occurs more quickly in downwind samples.


-UV transparent samples lose mass faster than the other treatments, especially when fertilized and downwind (site*bag*fertilization $P = 0.004$)



-Litter chemistry analyzed for 10 week samples:

-There is a lower C:P ratio in fertilized vs. non-fertilized ($P < 0.001$), particularly for downwind ($P < 0.001$) and UV opaque ($P = 0.003$)

-With regards to C:N ratios, downwind is lower than core ($P = 0.007$) and fertilized bags are marginally lower than control ($P = 0.06$)



Implications:

-This experiment shows that mass loss is more rapid in downwind samples.

-We determined how photodegradation vs. biological decomposition influences nutrient dynamics during litter degradation. Specifically, we discovered that allowing UV penetration allows for further degradation and lower C:P ratios.

Acknowledgements: Thanks go out to Jessica Alvarez Guevara, Jack Ramsey, and Bryan Beck for their help in the field and lab. This experiment wouldn't have been successful without their help. This study was funded by New College of Interdisciplinary Arts & Sciences.